

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 December 2003 (04.12.2003)

PCT

(10) International Publication Number
WO 03/099815 A1

(51) International Patent Classification⁷: **C07D 455/04**,
A61K 31/4375, A61P 31/04

(21) International Application Number: **PCT/IN02/00123**

(22) International Filing Date: **28 May 2002 (28.05.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(71) Applicant (for all designated States except US): **WOCKHARDT LIMITED [IN/IN]**; Wockhardt Towers, Bandra-Kurla Complex, Bandra (East), Mumbai 400 051 (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DE SOUZA, Noel, J. [IN/IN]**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN). **DESHPANDE, Prasad, K. [IN/IN]**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN). **SHUKLA, Milind, C. [IN/IN]**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN). **MUKARRAM, Siddiqui, M., Jaweed [IN/IN]**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN). **KULKARNI, Dilip, G. [IN/IN]**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN). **YEOLE, Ravindra, D. [IN/IN]**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN). **PATEL, Mahesh,**

V. [IN/IN]; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN). **GUPTE, Shrikant, V. [IN/IN]**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN).

(74) Agent: **DE SOUZA, Noel, J.**; Wockhardt Research Centre, D-4, MIDC Area, Chikalthana, Aurangabad 431 210, Maharashtra (IN).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **CRYSTALLINE FLUOROQUINOLONE ARGININE SALT FORM**

(57) Abstract: The invention relates to the new arginine salt forms of RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxy-1-piperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxy-1-piperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxy-1-piperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid, a process for their preparation and pharmaceutical formulations which comprise those arginine salt forms as the active ingredient for its use in treating microbial infections.

CRYSTALLINE FLUOROQUINOLONE ARGININE SALT FORM

FIELD OF THE INVENTION

The present invention relates to a new crystalline form of a fluoroquinolone, viz. arginine salt form thereof, a novel process for manufacturing the novel arginine salt form of the fluoroquinolone, the use of the novel form of the arginine salt of the fluoroquinolone in the manufacture of pharmaceutical formulations and the use of the novel form of the arginine salt of the fluoroquinolone in medicine. More particularly, it relates to the arginine salt form of the chiral fluoroquinolone S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid, a process for preparing the same and its use in pharmaceutical formulations and medicine.

BACKGROUND OF THE INVENTION

The chiral fluoroquinolone known under the name S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid is described in JP Patent 63,192,753A, JP Patent 05,339,238A, and in our pending U.S. Patent Applications Nos. 09/566,875, 09/640,947 and 09/802,793 and WO 00/68229, PCT Application Nos. PCT/IN00/00111 and PCT/IN01/00097.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid is an optically active isomer of the racemic compound which is claimed in U.S. Patent 4,399,134.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid has an aqueous solubility of 0.8 - 2.0 mg/ml over the pH range 8.0 - 9.5 at 28°C, thus creating problems in having to formulate the drug as a tablet or capsule, or in making formulations for gavage and parenteral injection. The need for a salt is clearly indicated, as the lack of an appropriate salt form can hinder the development of dosage forms acceptable for systemic use in mammals.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid has a pKa value of 6.80 suggesting a weak acid character and thus an ability to form a salt with an appropriate base. Generally, conversion of a pharmacologically active compound into a salt form induces a change in the compound's physicochemical properties such as solubility, absorption velocity, etc.

Pharmaceutically more desirable salt forms may be selected by studying whether or not a crystalline or amorphous form, or polymorph or pseudopolymorph can be produced, and determining the properties including its physicochemical or biological properties. A pseudopolymorph is a polymorph that differs from a true polymorph by the incorporation of solvent (Solid-state Chemistry of Drugs, 2nd Ed. S.R. Byrn et al (Eds). SSCI, Inc. 1999, p-514).

Pharmaceutically acceptable salts of racemic 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid such as salts with inorganic bases and organic bases are mentioned in the text of Otsuka's U.S. Patent No.4,399,134. Besides salts with inorganic bases and organic bases, amino acid salts of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid are identified in our pending U.S. Patent Applications No. 09/566,875 and U.S. Patent Application No. 09/640,947, WO 00/68229 and PCT Application No. PCT/IN00/00111. Arginine salt forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid are identified in our pending US Patent Application No. 09/802,793 and PCT Application No. PCT/IN01/00097. The subject matter of these applications is incorporated herein by reference. To date, no literature reference teaches about the amino acid salts of RS-(+/-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid or R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid, L-arginine salt 0.25 hydrate and S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid, L-arginine salt, 0.75 hydrate described in Examples 7 and 8 of U.S. Patent Applications 09/566,875 and 09/640,947, WO 00/68229 and PCT Application No. PCT/IN00/00111 respectively are highly

hygroscopic and turn into syrups on exposure at a relative humidity of 41%. Additionally, arginine salt forms described in our pending U.S. Patent Application No. 09/802,793 and PCT Application No. PCT/IN01/00097 are either amorphous or only partially crystalline and are to a certain degree hygroscopic. Many hydrates and salts associated with water are susceptible to changes in humidity, are hygroscopic under adverse storage conditions and during pharmaceutical processing of them to medicament forms.

The amino acid salts, inorganic base and alkali salts and organic base salts and specially the arginine salt forms of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo [i,j] quinolizine-2-carboxylic acid were prepared and studied by the inventors and it was found that:

- (a) an arginine salt may exist in a crystalline form having distinctive physicochemical, solubility and stability properties;
- (b) the crystalline arginine salt is less prone than the sodium salt to absorb moisture at specified humidity levels;
- (c) the crystalline arginine salt, possesses a lower propensity to cause phlebitis than the sodium and potassium salts as determined in rats by intravenous administration; and
- (d) the crystalline arginine salt is less toxic in rodents than the alkali salt forms.


In summary, the crystalline form of the arginine salt of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid has been found by the inventors to have very desirable properties in possessing under specified conditions, less hygroscopicity, favourable aqueous solubility, a low propensity to cause phlebitis, and favourable acute toxicity values. This form is expected to be very useful as a pharmaceutical agent as compared with the previously described arginine salt forms, sodium salt, other inorganic base/alkali salts, organic base salts and other amino acid salts. These advantages will be apparent from the experimental data shown hereafter.

SUMMARY OF THE INVENTION

New arginine salts of RS- (\pm)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid and S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid are described.

Compositions comprising one or more of these salts and methods for preparing the salts are described. The use of the salts in medicine is also described.

More particularly, a new L-arginine salt form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid of the formula I



Formula I

is described in which x denotes 0, 0.25, 0.5, 0.75, 1.0, 2.0 or 3.0. The novel crystalline form of the L-arginine salt of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid can be used in full scale manufacturing of pharmaceutical formulations.

Antibacterial compositions comprising the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt form as an active component are described.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention will now be described in further detail with reference to the accompanying drawings.

FIG. 1 represents the X-Ray Powder Diffraction (XRPD) spectrum of the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt of the invention.

FIG. 2 represents the Differential Scanning Calorimeter (DSC) thermogram of the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt of the invention.

FIG. 3 shows that the bulk density of the crystalline form of the invention S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt is between 0.57 g/cm³ (untapped) and 0.64 g/cm³ (tapped).

DISCLOSURE OF THE INVENTION


The main objective of the invention accordingly relates to new DL arginine salts and D-arginine salts and L-arginine salts of RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid and S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid.

Where particular stereoisomeric forms are described in this invention, they are meant to be substantially free of any other stereoisomeric configuration. Substantially free should be taken to mean that the active ingredient contains at least 90% by weight of the desired stereoisomer and 10% by weight or less of other stereoisomers. Preferably the weight % ratio is better than 95:5 and most preferably 99:1 or better. For example, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine contains at least 90% by weight of this stereoisomer and 10% by weight or less of other

configurations of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine.

The invention relates to new crystalline arginine salts of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid, wherein the arginine component may be L-arginine, D-arginine or DL-arginine.

More particularly, the invention relates to a new crystalline L-arginine salt of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid of the formula I



in which x denotes 0, 0.25, 0.5, 0.75, 1.0, 2.0 or 3.0. Preferably, the L-arginine crystalline salt is a monohydrate.

The salt of Formula I has favourable aqueous solubility, is stable in humid conditions, has favorable bioavailability, has a lower propensity to cause phlebitis and lower toxicity when administered in mammals.

This invention also relates to a process for the preparation of the salt of Formula I, to its use in pharmaceutical compositions and in medicine.

The arginine isomer used can be a racemic mixture of the chiral isomers or the R-isomer or the S-isomer. A preferred arginine isomer is the S-isomer, that is L-arginine.

The arginine salt of the invention is described as crystalline because of the high degree of crystallinity of the form as depicted by the XRD spectrum provided in Figure 1. The form is shown to have distinct physicochemical properties, is more stable to decomposition from uptake

of moisture at specified conditions, is distinguished by increased stability, in particular during storage at specified humidities and can be dried without caking or decomposing at elevated temperatures under reduced pressure. This form is particularly suitable for the preparation of stable pharmaceutical preparations, and has favourable biological properties.

This form, while less hygroscopic than the compounds of Examples 7 and 8 of U.S. Patent Applications 09/566,875 and 09/640,947, WO 00/68229 and PCT Application No. PCT/IN/00/00111, is to a certain degree hygroscopic. In stability studies at varying relative humidity conditions, this form was found to remain stable at relative humidity values of 22% at 25°C. In stability studies at elevated temperatures, this form remained stable with no decomposition up to 100°C, however, there was reduction in levels of water content and solvent content. On drying the forms at elevated temperatures up to 100°C over different periods of time and under different humidity conditions, the amounts of water found associated with the salt are altered. However, there is no change in the XRD-spectrum of the form. Without being bound to any theory, from this the inventors infer that the amount of water associated with the preferred form of the invention is not superficial and the preferred form of the invention is substantially a monohydrate.

Hydrates are said to exist when the drug substance incorporates water in the crystal lattice in either stoichiometric or non-stoichiometric amounts" (Stephen Byrn, et. al., Pharmaceutical Solids: A strategic Approach to Regulatory Considerations, Pharma. Res. Vol. 12 (7), 1995, 945-954).

The inventors have found that S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid arginine salt form of this invention, has acceptable aqueous solubility of 1 mg/ml extending to 30 mg/ml, over the pH range 7.0 - 9.5 at ambient temperatures. The solubility of this crystalline form is 30 mg/ml at pH 9.5. The L-arginine salt form is stable on heating at temperatures up to 100°C at relative humidity ranges up to 22%. This form has an acceptable dissolution rate.

The combination of physical properties of the novel S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-

arginine salt form of the present invention with respect to the degree of crystallinity, particle diameter, density, hygroscopicity, water content and content of other solvents is favourable and permits the manufacturing of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt in a composition which possesses the desired properties.

The properties of this crystalline form including the melting point is of a value that endow the form with desirable compression and flow properties for the processing of dosage forms useful for medicinal purposes.

This crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid arginine salt also has desirable physicochemical properties as hereinbefore mentioned, and is characterised by parameters such as XRD, DSC, particle size and powder density.

The more crystalline form is described below:

- (a) The degree of crystallinity as determined by X-ray powder diffraction is shown in Fig. 1, X-ray powder diffraction (2θ): 14.02 ± 0.2 , 14.82 ± 0.2 , 19.28 ± 0.2 , 22.12 ± 0.2 , 22.96 ± 0.2 , 23.46 ± 0.2 , 28.36 ± 0.2 .
- (b) A thermogram as determined by Differential Scanning Calorimetry has shown an exotherm at 194.93°C (onset at 189.42°C) and one endotherm at 251.26°C as shown in Fig. 2. Other endotherms at 87.83°C , 98.04°C , 144.03°C and 159.14°C are also observable, which endotherm readings may slightly fluctuate in different samples.

It is desirable that the crystalline form also has the following properties:

- (c) Particle size measured as mean mass diameter (MMD) as determined by laser diffraction technique is less than or equal to $40\mu\text{m}$, preferably the mean mass diameter is $32.00\mu\text{m} \pm 8\mu\text{m}$ as determined by laser diffraction technique, and as shown in Table 1;
- (d) Density between 0.57 g/cm^3 (untapped) and 0.64 g/cm^3 (tapped) as shown in Fig.3. Untapped density is defined as $25\text{ gm/untapped volume}$. The results are provided in Fig. 3. Tapped density is defined as $25\text{ gm/tapped volume}$. The results are provided in Fig. 3.
- (e) Hygroscopicity not exceeding 3% increase of weight upon storage for 14 days up to 22% relative atmospheric humidity at 30°C as determined gravimetrically, but with full retention

of flow properties;

- (f) A content of water between 4.0% to 4.4% by weight as determined by titration according to Karl Fischer.
- (g) The content of the organic solvent is less than 0.05% by weight as determined by gas chromatography. When acetone is used as solvent in the preparation of this form, the content of acetone is less than 0.3 %, preferably less than 0.2% by weight as determined by gas chromatography. When acetonitrile is used as solvent in the preparation of this form, the content of acetonitrile is less than 0.04 %, preferably less than 0.03 % by weight as determined by gas chromatography.
- (h) The solubility in a solution of pH 9.5 is 30.0 mg/ml.

The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt of this invention exhibits the same antibacterial minimum inhibitory concentration (MIC) values as the free active ingredient, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H- benzo [i,j] quinolizine-2-carboxylic acid.

The crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid L-arginine salt has antimicrobial activity:

It has potent activity against resistant *Staphylococcus aureus*.

It has ability to resist drug efflux in Gram-positive organisms.

It has the unusual ability to retain potency at an acidic pH of 5.5.

It is effective in treating respiratory pathogens.

It is effective in treating resistant mutants.

It has superior cidal action against slow-growing *Staphylococci*.

It scores over the standard drugs in terms of eradication efficacy for *Staphylococci* from vital organs and thigh muscle.

It exhibits unusual superior cidal effect even with a high density bacterial inoculum (10^8 cfu/ml); a standard recommended inoculum is 10^6 cfu/ml.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]-quinolizine-2-carboxylic acid D-arginine salt, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid DL-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid L-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid D-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid DL-arginine salt, R, S-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid L-arginine salt, R,S-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid D-arginine salt and R, S-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid DL-arginine salt also have antimicrobial activity. It is to be noted that S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid has more potent antimicrobial activity than R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid and RS(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]-quinolizine-2-carboxylic acid, this same difference in potency will also be displayed by their respective arginine salts.

The present invention also relates to a process for preparing the novel crystalline form of the arginine salt. A process for the manufacture of the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt comprises the following consecutive steps:

- a) dissolving S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid with L-arginine in an organic solvent and water to form a solution;
- b) cooling the solution to provide a crystalline substance;
- c) isolating the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4- hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt at < 35°C by

filtration or centrifugation;

- d) purifying and drying the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt using conventional methods.

The ratio of L-arginine to S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H, benzo [i,j] quinolizine-2-carboxylic acid ranges from 0.93 to 1.49 molar equivalent of L-arginine to 1.0 molar equivalent of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H, benzo [i,j] quinolizine-2-carboxylic acid. Preferably the molar ratio of L-arginine to S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H, benzo [i,j] quinolizine-2-carboxylic acid is 1.0:1.0 mole.

Preferably the solvents used in step (a) are an organic solvent and water; more preferably the organic solvent is acetone or acetonitrile. The proportion of organic solvent to water ranges from 20% to 80% depending upon the organic solvent used. The proportion of water to organic solvent ranges from 16.6% water to 33.3% water of the total volume of organic solvent and water used. A preferred proportion of water to organic solvent is 20% water of the total volume of water and organic solvent used.

Preferably the solution of step (a) is stirred at a temperature 50 to 60°C for 45-60 minutes.

Preferably in step (a) the solution is treated with activated charcoal at 50 to 60°C, filtered hot and diluted with an additional amount of organic solvent. In the case when the organic solvent is acetone, the ratio ranges from 0.9 to 1.1 parts by weight of the acid to 2.4 to 2.6 parts by volume of water to 9.0 to 11 parts by volume of acetone, preferably 1 part by weight of acid to 2.5 parts by volume of water to 10 parts by volume of acetone. In the case when the organic solvent is acetonitrile, the ratio ranges from 0.9 to 1.1 parts by weight of the acid to 2.3 to 2.5 parts by volume of water to 11.0 to 13 parts by volume of acetonitrile, preferably 1 part by weight of acid:2.4 parts by volume of water:12 parts by volume of acetonitrile.

The amount of water used depends on the nature of the organic solvent used. When the organic solvent is acetone, the preferred ratio of water to acetone is 1 part by volume of water to 4 parts

by volume of acetone and when acetonitrile is the organic solvent used, the preferred ratio of water to acetonitrile is 1 part by volume of water and 5 parts by volume of acetonitrile. It is also based on the amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H, benzo [i,j] quinolizine-2-carboxylic acid and the nature of the organic solvent used, preferably when 1 part by weight of acid is used, 2.5 parts by volume of water are used and 5 parts to 10 parts by volume of acetone are used. When acetonitrile is the solvent, preferably for 1 part by weight of acid, 2.4 parts by volume of water and 12 parts by volume of acetonitrile are used.

In step (b), the solution is cooled from 0°C to 40°C, more preferably from 30°C to 35°C.

Preferably in step (c), a crystalline substance is obtained by filtration.

In a preferred embodiment, L-arginine is added to a suspension or solution of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid in an organic solvent and water. The organic solvent may be selected from acetone or acetonitrile. The amounts of organic solvent and water used are so manipulated that when the reaction mixture is stirred for 45 – 60 minutes at 50 to 60°C a solution is provided, and subsequent cooling from 0°C to 40°C, preferably to 30 – 35°C provides the crystalline salt of the invention. The desired salt is isolated by filtration or centrifugation and washed with additional amount of the respective organic solvent used and dried preferably under reduced pressure and heating up to 65 – 70°C.

Yet a further aspect of the invention is that the synthesis of optically pure S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid arginine salt is also preferably prepared by enhancement of the optically impure or partially pure S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid using L-arginine, followed by crystallisation of the resulting optically pure salt. A less optically pure S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid used in this process may range from a mixture of 70% of S-(-)-isomer and 30% of R-(+)-isomer to 97% of S-(-)-isomer and 3% of R-(+)-isomer, preferably a mixture of 88% of S-(-)-isomer and 12% of R-(+)-isomer.

A general method for enhancement in optical purity of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid using L-arginine comprises,

(a) suspending a partially optically impure mixture of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i, j] quinolizine-2-carboxylic acid comprising 70% S-(-) isomer and 30% R-(+)- isomer to 97% S-(-) isomer and 3% R-(+)- isomer, preferably 88% S-(-) isomer and 12% R-(+)- isomer in water and organic solvent mixture. The ratio of water to organic solvent can be selected from 1.0:1.0 to 1.0:2.0, preferably 1.0:1.18. Organic solvent can be selected from acetone or acetonitrile, preferably acetone. An equimolar quantity of L-arginine is added to the suspension and the suspension is heated at 40 to 70°C, preferably 50 to 60°C to obtain a clear solution. 2 to 3 times more amount of the organic solvent than originally used is added, preferably 2.4 to 2.5 times more, and the suspension is stirred at 0 to 45°C, preferably 30 to 35°C for 1 hr to 5 h, preferably 2 hr, to effect the crystallization. The product is isolated by filtration and drying.

Yet a further aspect of the invention is that the synthesis of optically pure S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid arginine salt is also preferably prepared by enhancement of the optically impure or partially pure S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid using D-arginine, followed by crystallisation of the resulting optically pure salt. A less optically pure S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid used in this process may range from a mixture of 70% of S-(-)-isomer and 30% of R-(+)-isomer to 97% of S-(-)-isomer and 3% of R-(+)-isomer, preferably a mixture of 88% of S-(-)-isomer and 12% of R-(+)-isomer.

The process for manufacturing the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid L-arginine salt differs from the earlier known processes in that it can be controlled to give a crystalline polymorphic form of the salt form through use of the respective conditions that are described and shown to provide the result. This process can be carried out in conventional chemical process equipment. Preferably the product that is produced is substantially a

monohydrate.

Another aspect of the invention is the preparation of the DL- and D-arginine salt of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid and of the DL-, D- and L-arginine salts of R(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid and DL-, D- and L- arginine salt of RS(±)- 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid

The process comprises using a suspension of the appropriate enantiomer/racemic mixture of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid in a C₁-C₃ alkanol solvent, preferably methanol, ethanol, isopropanol, adding an aqueous solution of the appropriate enantiomer of arginine in preferably equimolar quantities, stirring the mixture at 40 – 70°C, preferably 60 – 65°C for 10 – 45 minutes, preferably 15 minutes, evaporating the solution to dryness under reduced pressure to give the desired salt.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid may be prepared by following the procedure described in U.S. Patent 4,399,134. L-arginine salt and D-arginine are commercially available from Ajinomoto, Japan. The cost of naturally occurring L-arginine is considerably cheaper than the enantiomeric D-arginine. One facet of this invention is the description of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine which utilizes the inexpensive naturally occurring and readily available L-arginine for making the crystalline salt of the invention.

A comparison between the novel form of the L-arginine salt prepared by the process of the present invention and those obtained from the experiments disclosed in the prior art shows that the salt of the present invention is substantially a monohydrate, is stable up to 100°C, and at relative humidity up to 22% at 25°C, and has favorable aqueous solubility, all attributes essential to the preparation of pharmaceutical compositions. This novel crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt is stable under typical storage conditions, has good bioavailability

in mammals, has lower phlebitis-forming potential on administration to mammals, has low or reduced toxicity, has acceptable disintegration and dissolution rates, and hence is very useful for pharmaceutical manufacturing and for use in medicine. The form is specially suitable for treating diseases caused by microbial infections. The form is suitable for long-term intravenous therapy in critically ill patients or patients in intensive care units. Injectable preparations of the L-arginine salt can be readily prepared in view of its availability in a bulk form that remains stable under specified conditions, its favorable aqueous solubility, its ideal suitability in not causing venous inflammation on repeated intravenous administration, and its safety from adverse toxicity.

S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt form of this invention can be present in pharmaceutical formulations as the only active compound or can be combined with other active ingredients such as other antibacterial agents.

The invention therefore also relates to liquid and solid pharmaceutical formulations which comprise the arginine salt of the invention, such as for example, injectable solutions, suspensions, emulsions, tablets, coated tablets, coated tablet cores, capsules, solutions, troches, dispersions, patches, powders, lotions, gels, sprays, pellets, granules, suppositories, hard or soft gelatin capsules, ointments, creams and the like.

The pharmaceutical formulations are prepared in a manner known per se, for example by mixing, stirring, suspending, dispersing, emulsifying, dissolving and the like, the active compounds with or in the pharmaceutical auxiliaries such as a carrier, diluent, solvent or excipient and processing the components to pharmaceutically suitable forms for parenteral, oral, topical, intranasal, buccal or rectal administration and the like.

Pharmaceutical formulations can be formulated together with auxiliaries and additives usually employed in pharmacy, such as tablet binders, fillers, preservatives, tablet disintegrating agents, flow regulating agents, plasticizers, wetting agents, dispersing agents, emulsifiers, solvents, pH altering additives, flavourings and the like.

The total daily dose range is generally from about 200 mg to about 1500 mg of the arginine salt

form. However, the dose may be higher or lower depending on the needs and conditions of the patient.

The following detailed examples serve to more fully illustrate the invention without limiting its scope.

EXAMPLE 1

Preparation of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt hydrate

S-(-)-9-Fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid (750 gm, 2.083 mole) was suspended in acetone (2250 ml). To this suspension was added L-arginine (337.5 g, 1.939 mole) and water (1875 ml) under stirring. The mixture was stirred at 55 to 60°C for one hour to obtain a clear solution. To the solution was added activated carbon (37.5 gm) and the solution was filtered hot. To the filtrate was added acetone (5250 ml). The reaction mixture was stirred for an additional for 2 hours at 30-35°C then allowed to cool to 5°C. The obtained solid was filtered at suction and washed with acetone. The wet solid was dried in a vacuum oven at 80-85°C to afford titled compound as a cream coloured powder. Yield 1035 gm. melting point 238-243°C. $[\alpha]_D^{25} = -184.25^\circ$ (c=1, methanol).

EXAMPLE 2

Preparation of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt hydrate

To a three-necked round bottom flask fitted on oil bath and equipped with magnetic stirrer and reflux condenser; S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid (2 gm, 5.55 mmoles) and L- arginine (1.44 gm, 8.27 mmole) were charged in a mixture of acetone (10 ml) and water (5 ml). The reaction mixture was slowly heated under stirring at 50 to 60°C temperature to obtain a clear solution. The solution was allowed to cool to 30 -35°C and then was further cooled in ice bath. The solid separated was filtered at suction and was washed with a cold mixture of acetone (10 ml) and water (5 ml). The crystalline solid was dried in a vacuum oven at 65-70°C to afford 2.4 gm (81%) dry compound, $[\alpha]_D^{25}$ value calculated on anhydrous basis is 183° (c= 1, methanol).

EXAMPLE 3

Preparation of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt hydrate

To the suspension of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid (100 gm, 0.278 mole) in acetone (300 ml) was added L- arginine (45 gm, 0.258 mole) and water (250 ml). The mixture was heated to 50 to 60°C temperature for one hour to obtain a clear solution. Activated charcoal (5 gm) was added to the solution and the solution was filtered hot. To the filtrate was added acetone (700 ml) and the mixture was cooled gradually to 30 to 35°C temperature, a solid crystallized out which was filtered and washed with aqueous acetone and finally with acetone. On drying in vacuum oven at 65 to 70°C S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid, L-arginine salt was obtained as a crystalline powder weighing 137 gm (92%). $[\alpha]_D^{25}$ value calculated on anhydrous basis is -183.34° (c=1, methanol)

EXAMPLE 4

Preparation of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt hydrate

To a three necked 1 lit. round bottom flask fitted on the water bath and equipped with mechanical stirrer, thermometer pocket and reflux condenser was charged S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid (25 gm, 69.44 mmole) and acetonitrile (75 ml). Stirring was started and to the stirred suspension was charged powdered L- arginine (12.16 gm, 69.88 mmole) followed by distilled water (60 ml). The reaction mixture was stirred at the temperature between 50 to 60°C for one hour to obtain a homogeneous clear solution. Clear reaction mixture was then charged 225 ml fresh acetonitrile under stirring. Activated charcoal (3 gm) was added to the solution and the solution was filtered hot. Filtrate was allowed to cool to 30 to 35°C under stirring. The reaction mixture was stirred for additional 3 hours at this temperature. The crystalline solid was filtered

at suction and the wet cake was washed with 25 ml acetonitrile. Resulting crystalline solid was dried under vacuum at 65 to 70°C to furnish 34.5 gm (93%) title compound, $[\alpha]_D^{25}$ value calculated on anhydrous basis is -183.99° ($c=1$, methanol).

EXAMPLE 4-A

Preparation of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt

A mixture of the two enantiomers [88.24% S-(-)-isomer and 11.76% R-(+)-isomer] of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid (1.5 gm, 4.16 mmol) was suspended in a mixture of acetone (4.5 ml) and water (3.8 ml). To the stirred suspension L-arginine (0.725 gm, 4.16 mmol) was added. The mixture was warmed to a temperature between 50-60°C to obtain a clear solution. Acetone (11 ml) was added to the solution, which was then cooled to 30 - 35 °C and stirred for 2 hours. The crystalline solid obtained was filtered under suction and the residue was washed with 2 ml acetone. The resulting solid was dried to furnish 1.8 gm (92%) of the title compound.

Chiral analysis of the crystals were performed by chromatography on an analytical chiral column. The optical purity of the crystals was found to be 97.06%. The crystallinity was determined by powder X-ray diffraction.

EXAMPLE 4-B

The above experiment was repeated on the mixture of enantiomer having optical purity 89% S-(-)-isomer and 11% R-(+)-isomer of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo 1H,5H-benzo[i,j] quinolizine-2-carboxylic acid using acetonitrile instead of acetone in accordance with the conditions and quantities described in Example 4-A to provide the product of which optical purity was enhanced to 98% as shown by chiral chromatography.

EXAMPLE 5

Preparation of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzof[i,j]quinolizine-2-carboxylic acid D-arginine salt

To a suspension of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid (1 g, 2.77 mmol) in methanol (50 ml) was charged a solution of D-arginine (0.483 g, 2.77 mmol) in 10 ml water at 60 – 65°C. The clear solution obtained was stirred for 15 minutes at 60 – 65°C. The reaction mixture was evaporated to dryness under reduced pressure to give 1.3 g (88%) of the title compound, m.p. 241 – 243°C, $[\alpha]_D^{25} = -174.4^\circ$ (c=1, methanol).

EXAMPLE 6

Preparation of R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt

This compound may be prepared according to the procedure described in Example 5 using R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid instead of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid and L-arginine instead of D-arginine.

EXAMPLE 7

Preparation of R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt

This compound may be prepared following Example 6 by substituting an equivalent amount of D-arginine for L-arginine. m.p. (DSC) 232.0°C, $[\alpha]_D^{25} = +171.4^\circ$ (c=1, methanol)

EXAMPLE 8

Preparation of RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt

This compound may be prepared following Example 6 by substituting RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid for R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid.

EXAMPLE 9

Preparation of RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt

This compound may be prepared following Example 8 by substituting an equivalent amount of D-arginine for L-arginine. M.p (DSC) 234.8°C.

The crystalline S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt prepared according to Examples 1-4B possesses the following properties:

- a) Crystalline form, with a degree of crystallinity as determined by X-ray powder diffraction and as shown in Fig. 1.
- b) A thermogram as determined by Differential scanning calorimetry and as shown in Fig. 2.
- c) Particle size measured as mean mass diameter (MMD) of less than or equal to 40 µm, as determined by laser diffraction technique.
- d) Bulk density of 0.57 g/cm³ (untapped) and 0.64 g/cm³ (tapped) as shown in Fig 3. Untapped density is defined as 25 gm/untapped volume. The results are provided in Fig. 3. Tapped density is defined as 25 gm/tapped volume. The results are provided in Fig. 3.
- e) Hygroscopicity of 3% increase of weight upon storage for 14 days up to 22% relative atmospheric humidity at 30°C as determined gravimetrically.
- f) Water content of between 4.0% to 4.4% by weight as determined by titration according to Karl Fischer.
- g) A content of acetone of 0.011% by weight, or of acetonitrile of 0.03% by weight, as determined by gas chromatography.

TEST EXAMPLE 1

X-ray Diffraction Analysis

300 mg each of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [I,j]quinolizine-2-carboxylic acid L-arginine salt prepared as in Example 1 was thinly

spread on a sample holder. X-ray diffraction analyses (40kv x 40 mA Rigaku D/max 2200) were performed under the conditions listed below:

Scan speed 5°/ min

Sampling time 7 min

Scan mode: continuous

2 θ / θ reflection

Cu target (Ni filter)

Results of the X-ray diffraction analysis on the crystalline form is depicted in Fig. 1.

TEST EXAMPLE 2

Thermal Analysis of the S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [L,i]quinolizine-2-carboxylic acid L- arginine salt

For the Differential Scanning Calorimetry, METTLER TOLEDO STAR system was used. 5 to 6 mg of the sample was weighed into the aluminum pan, which was then press sealed with an aluminium lid. After three tiny needle holes were made on the lid the sample was tested by heating from (30°C) to (300°C) at a rate of 10°C/min. Result can be seen from Fig. 2. There is an exothermic peak at around 194.93°C and an endothermic peak at 251.26°C, other endothermic peaks are observable at 87.83°C, 98.04°C, 144.03°C and 159.14°C

TEST EXAMPLE 3

BULK DENSITY DETERMINATION

For untapped and tapped density determination Bulk Density Apparatus (R.V. Electronics, Mumbai) was used. 25 g of the sample was slowly poured into a dry clean stoppered measuring cylinder and filled volume was measured to obtain untapped density.

Untapped density = 25 g/untapped volume.

Then the measuring cylinder was fixed to Bulk Density Apparatus and sample was tapped (150 times) and volume was measured to furnish the tapped volume.

Tapped density = 25 g/Tapped volume.

The results are shown in Fig. 3.

TEST EXAMPLE 4
MEAN MASS DIAMETER DETERMINATION

For mean mass diameter Master Sizer of Malvern Instrument LTD U.K. was used under the conditions listed below:

Lens/Focus: 300 mm

Solvent: Hexane

Analysis model: Polydisperse

Obstruction value: 20-23%

The particle size measured as mean mass diameter of the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt of the invention from four different samples made by the procedure described in example 1 is as given in Table 1.


Table 1

Sample No.	Mean mass diameter (μm)
1	32.92
2	24.22
3	34.51
4	39.07

The mean value is 32.68 μm .

CLAIMS

1. A compound which is selected from the DL-, D- or L- arginine salts of racemic or optically active isomers of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid.
2. A compound according to claim 1, which is selected from the DL-, D- or L- arginine salts of optically active isomers of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid, which selected compound is substantially free of any other 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid – arginine salt stereoisomer.
3. A compound of claim 2, which is selected from S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt and S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid D-arginine salt.
4. A compound of claim 2, wherein the weight ratio of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt or S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt to other stereoisomers of this salt is at least 95:5.
5. A compound of claim 4, wherein the weight ratio of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt or S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid D-arginine salt to other stereoisomers of this salt is at least 99:1.
6. S-(-)-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt of the formula I



Formula I

in which x denotes 0, 0.25, 0.5, 0.75, 1.0, 2.0 or 3.0 in a crystalline form.

7. A S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 6, wherein $x = 1$.
8. A S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid L-arginine salt having the following X-ray powder diffraction data: (2θ) : 14.02 ± 0.2 , 14.82 ± 0.2 , 19.28 ± 0.2 , 22.12 ± 0.2 , 22.96 ± 0.2 , 23.46 ± 0.2 , 28.36 ± 0.2 .
9. A S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid L-arginine salt having the following X-ray powder diffraction data: (2θ) : 14.02 ± 0.2 , 14.82 ± 0.2 , 19.28 ± 0.2 , 22.12 ± 0.2 , 22.96 ± 0.2 , 23.46 ± 0.2 , 28.36 ± 0.2 ; a DSC exotherm at 194.93°C (onset at 189.42°C) and one endotherm at 251.26°C and a water content of between 4.0 and 4.4% by weight as determined by titration according to Karl Fischer.
10. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 8, having an DSC exotherm at 194.93°C (onset at 189.42°C) and one endotherm at 251.26°C .
11. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 8, wherein the solubility in a solution of pH 9.5 is 30.0 mg/ml.

12. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 9, wherein the solubility in a solution of pH 9.5 is 30.0 mg/ml.
13. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 8, wherein the mean mass diameter as determined by laser diffraction technique is less than or equal to 40 μm .
14. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 13, wherein the diameter is less than 32 μm .
15. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 8, wherein the density is between 0.57 g/cm³ (untapped) and 0.64 g/cm³ (tapped).
16. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 9, wherein the density is between 0.57 g/cm³ (untapped) and 0.64 g/cm³ (tapped).
17. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 6, wherein the water content is between 4.0 % and 4.4 % by weight as determined by titration according to Karl Fischer.
18. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 8, wherein the water content is between 4.0 and 4.4% by weight as determined by titration according to Karl Fischer.
19. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo

- [i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 8, having an organic solvent content less than 0.05 % by weight as determined by gas chromatography.
20. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 9, having an organic solvent content less than 0.05 % by weight as determined by gas chromatography.
21. The S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j] quinolizine-2-carboxylic acid L-arginine salt according to claim 6, wherein the hygroscopicity is less than 3 % increase of weight upon storage for 14 days at up to 22% relative atmospheric humidity as determined by gravimetrically.
22. S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid D-arginine salt having m.p. 241 – 243°C, and $[\alpha]_D^{25} = -174.4^\circ$ (c=1, methanol).
23. R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid L-arginine salt .
24. R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid D-arginine salt having m.p. (DSC) 232.0°C, and $[\alpha]_D^{25} = +171.4^\circ$ (c=1, methanol)
25. R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid DL-arginine salt .
26. A compound selected from RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid DL-arginine salt , RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid L-arginine salt or RS-(±)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i,j]quinolizine-2-carboxylic acid D-arginine salt.

27. A process for the manufacture of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt comprising the steps of:
- dissolving S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid with L-arginine in an organic solvent and water to form a solution;
 - cooling the solution to provide a crystalline substance;
 - isolating the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H, 5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt at < 35°C by filtration or centrifugation; and
 - purifying and drying the crystalline form of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt.
28. The according to claim 27, wherein the molar ratio of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid to L-arginine is in the range from 1.0: 0.93 to 1.49.
29. The process according to claim 27, wherein the molar ratio of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid to L-arginine is 1:1.
30. The process according to claim 27, wherein the organic solvent is selected from acetone or acetonitrile.
31. The process according to claim 27, wherein the percentage of water in the solvent mixture is 20% - 80%.
32. The process according to claim 27, wherein the solution of step a) is stirred at 50 to 60°C.

33. A method for enhancing optical purity of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j] quinolizine-2-carboxylic acid L-arginine salt comprising,
- (a) suspending a partially optically impure mixture of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo [i, j] quinolizine-2-carboxylic acid comprising 70% S-(-) isomer and 30% R-(+)- isomer to 97% S-(-) isomer and 3% R-(+)- isomer in water and organic solvent to form a suspension,
 - (b) adding adding an equimolar quantity of L-arginine to the suspension and heating suspension at 40 to 70°C to obtain a clear solution, and
 - (c) adding 2 to 3 times more of the organic solvent added in step a) and stirring at 0 to 45°C, for 1 hr to 5 hr, to effect the crystallization and isolating the product by filtration and drying.
34. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 6 and a carrier, diluent, solvent or excipient.
35. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 7 and a carrier, diluent, solvent or excipient.
36. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 8 and a carrier, diluent, solvent or excipient.
37. A composition comprising S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 9 and a carrier, diluent, solvent or excipient.
38. A composition comprising a compound selected from the group consisting of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-

methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt, RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt, RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt and RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt and a carrier, diluent, solvent or excipient.

39. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 6 to the mammal in need thereof.
40. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 7, to the mammal in need thereof.
41. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 8, to the mammal in need thereof.
42. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt according to claim 9, to the mammal in need thereof.

43. A method for treating a disease caused by a microbial infection in a mammal comprising administering an effective amount of a compound selected of the group consisting of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt, RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt, RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt and RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt to the mammal in need thereof.
44. A process for the manufacture of a compound selected from the group selected from the group consisting of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt, S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt, R-(+)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt, RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid L-arginine salt, RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt and RS-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid DL-arginine salt S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo[i,j]quinolizine-2-carboxylic acid D-arginine salt comprising the steps of:

- a) suspending an enantiomer or racemic mixture of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-oxo-1H,5H-benzo[i,j]quinolizine-carboxylic acid in a C₁-C₃ alkanol solvent and adding an aqueous solution of DL, L or D arginine;
 - b) stirring the mixture at 40-70°C for 10-45 minutes; and evaporating the solution to dryness under reduced pressure.
45. A process according to claim 44, wherein the amounts used of S-(-)-9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-oxo-1H,5H-benzo[i,j]quinolizine-carboxylic acid and D-arginine are equal molar amounts.
46. A process according to claim 44, wherein the solvent is selected from methanol, ethanol, or isopropanol.




FIG. 2





FIG. 3



INTERNATIONAL SEARCH REPORT

International Application No
PCT/IN 02/00123

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D455/04 A61K31/4375 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 68229 A (AGARWAL S K; CHUGH Y (IN); GUPTE S V (IN); PATEL M V (IN); SAOJI D) 16 November 2000 (2000-11-16) cited in the application Page 66, lines 15-22: (S)-Nadifloxacin-Arg salt, Arg salt 0.25 H2O, Arg salt 0.75 H2O; page 43: (S)-Nadifloxacin-L-Arg salt 0.25 H2O and 0.75 H2O; page 1; page 7 item 2; page 9 item 12.	1-46
X	WO 01 85095 A (PATEL MAHESH VITHALBHAI; YEOLE RAVINDRA DATTATRAYA (IN); SHETTY NI) 15 November 2001 (2001-11-15) Claims 1-28 and examples 1-5: (S)-Nadifloxacin-L-Arg salts; claims 44-55.	1-46



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

23 August 2002

Date of mailing of the international search report

02/09/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Weisbrod, T

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IN 02/00123

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02 09758 A (KHORAKIWALA HABIL F; NAIR SHEELA C (IN); WOCKHARDT LTD (IN); PATEL) 7 February 2002 (2002-02-07) Page 61, example 54: (S)-Nadifloxacin-L-Arg salt; table 1, page 91, entry 4, and claim 38, page 123, lines 49-50: (R)-Nadifloxacin-L-Arg salt. <u> </u>	1-46

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IN 02/00123

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 39-43 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound(s).
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1, 2, 38, 43, 44, 46 (all part) and 25, 26 (complete)

directed to DL-arginine (arginine = Arg) salts of racemic or optically active isomers of 9-fluoro-6,7-dihydro-8-(4-hydroxypiperidin-1-yl)-5-methyl-1-oxo-1H,5H-benzo(1,j)quinolizin-2-carboxylic acid (Nadifloxacin), a composition comprising such salts, a therapeutic method involving such salts, and a process for their manufacture,

2. Claims: 1-5, 26, 38, 43, 44, 46 (all part) and 22, 24, 45 (complete)

directed to D-Arg salts of racemic or optically active isomers of Nadifloxacin, a composition comprising such salts, a therapeutic method involving such salts, and a process for their manufacture,

3. Claims: 1, 26, 38, 43, 44, 46 (all part)

directed to L-Arg (rac)-Nadifloxacin salt, a composition comprising such salt, a therapeutic method involving such salt, and a process for its manufacture,

4. Claims: 1-6, 21, 27-32, 34, 39 (all part)

directed to the L-Arg (S)-Nadifloxacin salt respectively the salt L-Arg (S)-Nadifloxacin-xH₂O with x = 0, a composition comprising such salt, a therapeutic method involving such salt, and a process for its manufacture,

5. Claims: 1, 2, 38, 43, 44, 46 (all part) and 23 (complete)

directed to L-Arg (R)-Nadifloxacin salt, a composition comprising such salt, a therapeutic method involving such salt, and a process for its manufacture,

6. Claims: 6, 21, 27-32, 34, 39 (all part)

directed to L-Arg (S)-Nadifloxacin-xH₂O with x = 0.25, a composition comprising such salt, a therapeutic method involving such salt, and a process for its manufacture,

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

7. Claims: 6, 21, 27-32, 34, 39 (all part)

directed to L-Arg (S)-Nadifloxacin-xH₂O with $x = 0.5$, a composition comprising such salt, a therapeutic method involving such salt, and a process for its manufacture,

8. Claims: 6, 21, 27-32, 34, 39 (all part)

directed to L-Arg (S)-Nadifloxacin-xH₂O with $x = 0.75$, a composition comprising such salt, a therapeutic method involving such salt, and a process for its manufacture,

9. Claims: 6, 21, 27-32, 34, 39 (all part) and 7-20, 35-37, 40-42 (complete)

directed to L-Arg (S)-Nadifloxacin-xH₂O with $x = 1.0$, L-Arg (S)-Nadifloxacin salt with a water content of 4.0-4.4% (corresponding with L-Arg (S)-Nadifloxacin-xH₂O wherein x is around 1.2), as well as L-Arg (S)-Nadifloxacin salt as defined by its X-ray diffraction data according to claims 8 and 9, a composition comprising such salt(s), a therapeutic method involving such salt(s), and a process for the manufacture of such salt(s),

10. Claims: 6, 21, 27-32, 34, 39 (all part)

directed to L-Arg (S)-Nadifloxacin-xH₂O with $x = 2.0$ or 3.0 , a composition comprising such salts, a therapeutic method involving such salts, and a process for the manufacture of such salt(s),

11. Claim : 33 (complete)

directed to a method for enhancing the optical purity of L-Arg (S)-Nadifloxacin salt.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/IN 02/00123

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0068229	A	16-11-2000	AU 5425600 A 21-11-2000
		EP 1175217 A2 30-01-2002	
		WO 0068229 A2 16-11-2000	
WO 0185095	A	15-11-2001	AU 7866601 A 20-11-2001
		AU 8009101 A 13-02-2002	
		WO 0185095 A2 15-11-2001	
		WO 0209758 A2 07-02-2002	
		US 2002061908 A1 23-05-2002	
		AU 7866701 A 20-11-2001	
		WO 0185728 A2 15-11-2001	
WO 0209758	A	07-02-2002	AU 7866601 A 20-11-2001
		AU 8009101 A 13-02-2002	
		WO 0185095 A2 15-11-2001	
		WO 0209758 A2 07-02-2002	
		US 2002061908 A1 23-05-2002	
		AU 7866701 A 20-11-2001	